

## UTILIZATION MANAGEMENT MEDICAL POLICY

**POLICY:** Oncology (Injectable) – Proleukin Utilization Management Medical Policy

- Proleukin® (aldesleukin intravenous infusion – Prometheus Laboratories)

**REVIEW DATE:** 01/18/2023

---

### OVERVIEW

Proleukin, a human recombinant interleukin-2 product, is indicated for the following conditions, in adults:

- **Metastatic melanoma.**
- **Metastatic renal cell carcinoma.**<sup>1</sup>

### Dosing Information

The recommended dose of Proleukin is the same for metastatic melanoma and metastatic renal cell carcinoma.<sup>1</sup> Proleukin 600,000 International Units/kg (0.037 mg/kg) is administered by intravenous infusion over 15 minutes every 8 hours for a maximum of 14 doses. Following 9 days of rest the schedule is repeated to complete one course of therapy. Additional courses of therapy can be given after at least 7 weeks of rest. Additional courses of therapy should only be given if there is evidence of tumor shrinkage after the previous course of therapy and there are no contraindications to retreatment.

### Guidelines

Proleukin is addressed in the following National Comprehensive Cancer Network guidelines:

- **Cutaneous melanoma** (version 1.2023 – December 22, 2022) clinical practice guidelines recommend Proleukin for unresectable or metastatic disease as a single agent for second-line or subsequent therapy for disease progression or after maximum clinical benefit from BRAF targeted therapy (category 2A).<sup>2,4</sup> Proleukin may be considered for patients with small brain tumors and without significant peritumoral edema (category 2B) or for intralesional therapy as primary or second-line treatment of unresectable stage III disease with clinical or satellite/in-transit metastases, or local satellite/in-transit recurrence (category 2B).
- **Hematopoietic cell transplantation** (version 2.2022 – September 28, 2022) clinical practice guidelines recommend Proleukin as additional therapy, in combination with systemic corticosteroids, for steroid-refractory chronic graft-vs-host disease.<sup>2,5</sup>
- **Kidney cancer** (version 3.2023 – September 22, 2022) clinical practice guidelines recommend Proleukin as a single agent for first-line (category 2B) and subsequent (category 2B) therapy for patients with relapsed or stage IV disease and clear cell histology.<sup>2,3</sup>

### POLICY STATEMENT

Prior Authorization is recommended for medical benefit coverage of Proleukin. Approval is recommended for those who meet the **Criteria** and **Dosing** for the listed indications. Extended approvals are allowed if the patient continues to meet the Criteria and Dosing. Requests for doses outside of the established dosing documented in this policy will be considered on a case-by-case basis by a clinician (i.e., Medical Director or Pharmacist). All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Proleukin as well as the monitoring required for adverse events and long-term efficacy, approval requires Proleukin to be prescribed by or in consultation with a physician who specializes in the condition being treated.

**Automation:** None.



## RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Proleukin is recommended in those who meet one of the following criteria:

### FDA-Approved Indications

---

**1. Cutaneous Melanoma.** Approve for 1 year if the patient meets ONE of the following (A or B):

**A) Intravenous Therapy.** Approve if the patient meets the following criteria (i, ii, iii, iv, and v):

- i. Patient is  $\geq 18$  years of age; AND
- ii. Patient has metastatic or unresectable disease; AND
- iii. Patient has tried at least one other systemic therapy; AND
- iv. Proleukin will be used as a single agent; AND
- v. Proleukin is prescribed by or in consultation with an oncologist.

**B) Intralesional Therapy.** Approve if the patient meets the following criteria (i, ii, and iii):

- i. Patient is  $\geq 18$  years of age; AND
- ii. Proleukin will be directly injected into metastatic, recurrent, or unresectable cutaneous, subcutaneous, or nodal lesions; AND
- iii. The medication is prescribed by or in consultation with an oncologist or dermatologist.

**Dosing.** Approve one of the following dosing regimens (A or B):

**A) Intravenous Therapy** (i, ii, and iii):

- i. Each dose must not exceed 600,000 International Units/kg (0.037 mg/kg) given no more frequently than three times daily for a maximum of 14 doses to complete one cycle of treatment; AND
- ii. A second cycle is given after a minimum of 9 days of rest to complete a course of therapy; AND
- iii. Each additional course of therapy is given after at least 7 weeks of rest; OR

**B) Intralesional Therapy** (i and ii):

- i. The dose to each individual lesion must not exceed 6 million International Units given by intralesional injection; AND
- ii. The dose is given no more frequently than three times weekly.

---

**2. Kidney Cancer.** Approve for 1 year if the patient meets the following criteria (A, B, C, D, and E):

- A) Patient is  $\geq 18$  years of age; AND
- B) Patient has relapsed or metastatic disease; AND
- C) Patient has clear cell histology; AND
- D) Proleukin will be used as a single agent; AND
- E) Proleukin is prescribed by or in consultation with an oncologist.

**Dosing.** Approve the following dosing regimen (A, B, and C):

- A) Each dose must not exceed 600,000 International Units/kg (0.037 mg/kg) given intravenously no more frequently than three times daily for a maximum of 14 doses to complete one cycle of treatment; AND
- B) A second cycle is given after a minimum of 9 days of rest to complete a course of therapy; AND
- C) Each additional course of therapy is given after at least 7 weeks of rest.

### Other Uses with Supportive Evidence



**3. Graft-Versus-Host Disease.** Approve for 1 year if the patient meets the following criteria (A, B, C, and D):

- A) Patient has chronic graft-versus-host disease; AND
- B) According to the prescriber, the patient has steroid-refractory disease; AND
- C) Proleukin will be used in combination with systemic corticosteroids; AND
- D) Proleukin will be prescribed by or in consultation with an oncologist or a physician associated with a transplant center.

**Dosing.** Approve up to 1 million International Units/m<sup>2</sup> administered subcutaneously no more frequently than once daily.

---

**CONDITIONS NOT RECOMMENDED FOR APPROVAL**

Coverage of Proleukin is not recommended in the following situations:

- 1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

**REFERENCES**

- 1. Proleukin® intravenous infusion [prescribing information]. San Diego, CA: Prometheus Laboratories; September 2019.
- 2. The NCCN Drugs and Biologics Compendium. © 2023 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on January 10, 2023. Search term: aldesleukin.
- 3. The NCCN Kidney Cancer Clinical Practice Guidelines (version 3.2023 – September 22, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed January 10, 2023.
- 4. The NCCN Cutaneous Melanoma Clinical Practice Guidelines (version 1.2023 – December 22, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed January 10, 2023.
- 5. The NCCN Hematopoietic Cell Transplantation Clinical Practice Guidelines (version 2.2022 – September 28, 2022). © 2022 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed January 10, 2023.
- 6. Radny P, Caroli UM, Bauer J, et al. Phase II trial of intralesional therapy with interleukin-2 in soft-tissue melanoma metastases. *Br J Cancer*. 2003;89:1620-1626.
- 7. Weide B, Derhovanessian E, Pflugfelder A, et al. High response rate after intratumoral treatment with interleukin-2. Results from a Phase 2 study in 51 patients with metastasized melanoma. *Cancer*. 2010;116:4139-4146.
- 8. Koreth J, Kim HT, Jones KT, et al. Efficacy, durability, and response predictors of low-dose interleukin-2 therapy for chronic graft-versus-host disease. *Blood*. 2016;128:130-137.

**HISTORY**

Type of Revision	Summary of Changes	Review Date
Annual Revision	<b>Cutaneous Melanoma:</b> Increased approval duration to 1 year. <b>Kidney Cancer:</b> Increased approval duration to 1 year. <b>Graft-Versus-Host Disease:</b> Revised dose from International Units to International Units/m <sup>2</sup> .	12/22/2021
Annual Revision	No criteria changes.	01/18/2023