



## PRIOR AUTHORIZATION POLICY

- POLICY:** Complement Inhibitors – PiaSky Prior Authorization Policy
- PiaSky® (crovalimab-akkz intravenous infusion or subcutaneous injection – Genentech/Roche)

**REVIEW DATE:** 07/10/2024

---

### OVERVIEW

PiaSky, a complement C5 inhibitor, is indicated for the treatment of **paroxysmal nocturnal hemoglobinuria (PNH)** in patients  $\geq 13$  years of age who weigh  $\geq 40$  kg.<sup>1</sup>

### Disease Overview

Paroxysmal nocturnal hemoglobinuria (PNH) is a rare, genetic disorder of hematopoietic stem cells.<sup>2,3</sup> The mutation in the X-linked gene phosphatidylinositol glycan class A (PIGA) results in a deficiency in the glycosylphosphatidylinositol (GPI) protein, which is responsible for anchoring other protein moieties to the surface of the erythrocytes. Loss of anchoring of these proteins causes cells to hemolyze and leads to complications such as hemolytic anemia, thrombosis, and peripheral blood cytopenias. PNH is a clinical diagnosis that should be confirmed with peripheral blood flow cytometry to detect the absence or severe deficiency of GPI-anchored proteins on at least two lineages.<sup>2,5</sup> Prior to the availability of complement inhibitors, only supportive management, in terms of managing the cytopenias and controlling thrombotic risk were available. Supportive measures include platelet transfusion, immunosuppressive therapy for patients with bone marrow failure, use of erythropoietin for anemias, and aggressive anticoagulation.

### Dosing Information

The recommended dosage regimen for PiaSky consists of one loading dose administered by intravenous infusion on Day 1, followed by four weekly loading doses administered by subcutaneous (SC) injection on Days 2, 8, 15, and 22.<sup>1</sup> Maintenance doses, which are given once every 4 weeks by SC injection, start on Day 29. Only healthcare providers should administer PiaSky.

### Safety

The PiaSky prescribing information has a Boxed Warning about serious meningococcal infections.<sup>1</sup> PiaSky is only available through a restricted access program, PiaSky Risk Evaluation and Mitigation Strategy (REMS).

### POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of PiaSky intravenous. All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days. Because of the specialized skills required for evaluation and diagnosis of patients treated with PiaSky intravenous as well as the monitoring required for adverse events and long-term efficacy, approval requires PiaSky intravenous to be prescribed by or in consultation with a physician who specializes in the condition being treated.

**Automation:** None.

---

## RECOMMENDED AUTHORIZATION CRITERIA

Coverage of PiaSky is recommended in those who meet the following criteria:

### FDA-Approved Indication

1. **Paroxysmal Nocturnal Hemoglobinuria.** Approve for the duration noted if the patient meets ONE of the following (A or B):
  - A) **Initial therapy.** Approve for 6 months if the patient meets ALL of the following (i, ii, iii, and iv):
    - i. Patient is  $\geq 13$  years of age; AND
    - ii. Patient weighs  $\geq 40$  kg; AND
    - iii. Diagnosis was confirmed by peripheral blood flow cytometry results showing the absence or deficiency of glycosylphosphatidylinositol (GPI)-anchored proteins on at least two cell lineages; AND
    - iv. The medication is prescribed by or in consultation with a hematologist.
  - B) **Patient is Currently Receiving PiaSky subcutaneous.** Approve for 1 year if the patient meets ALL of the following (i, ii, iii, and iv):

Note: A patient who has not started maintenance therapy with PiaSky subcutaneous should be considered under criterion A (Initial Therapy).

    - i. Patient is  $\geq 13$  years of age; AND
    - ii. Patient weighs  $\geq 40$  kg; AND
    - iii. According to the prescriber, patient is continuing to derive benefit from PiaSky; AND  
Note: Examples of benefit include increase in or stabilization of hemoglobin levels, decreased transfusion requirements or transfusion independence, reductions in hemolysis.
    - iv. The medication is prescribed by or in consultation with a hematologist.

### CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of PiaSky is not recommended in the following situations:

1. **Concomitant Use with Another Complement Inhibitor.** There is no evidence to support concomitant use of PiaSky with another complement inhibitor.

Note: Examples of complement inhibitors are Empaveli (pegcetacoplan subcutaneous injection), Fabhalta (iptacopan capsule), Soliris (eculizumab intravenous infusion), Ultomiris (ravulizumab cwzy intravenous infusion or subcutaneous injection), Voydeya (danicopan tablets).
2. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

### REFERENCES

1. PiaSky<sup>®</sup> [prescribing information]. South San Francisco, CA: Genentech; June 2024.
  2. Cançado RD, da Silva Araújo A, Sandes AF, et al. Consensus statement for diagnosis and treatment of paroxysmal nocturnal haemoglobinuria. *Hematol Transfus Cell Ther.* 2021;43:341-348.
  3. Shah N, Bhatt H. Paroxysmal Nocturnal Hemoglobinuria. [Updated 2023 Jul 31]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023 Jan. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK562292/>. Accessed on July 1, 2024.
  4. Roth A, Maciejewski J, Nishinura JI, et al. Screening and diagnostic clinical algorithm for paroxysmal nocturnal hemoglobinuria: Expert consensus. *Eur J Haematol.* 2018;101(1):3-11.
-

**HISTORY**

<b>Type of Revision</b>	<b>Summary of Changes</b>	<b>Review Date</b>
New Policy	--	07/10/2024